

Purpose

- controls the symptoms of diarrhea, including Traveler's Diarrhea

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Do not use if you have bloody or black stool

Ask a doctor before use if you have

- high fever (greater than 101°)
- mucus present in your stool
- a history of liver disease

Ask doctor or pharmacist before use if you are taking antibiotics

Stop use and ask a doctor if diarrhea lasts for more than 2 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Do not use

Ask a doctor before use if you have

Ask a doctor or pharmacist before use if you are

When using this product

Stop use and ask a doctor if

If pregnant or breast-feeding

Keep out of reach of children

- use enclosed dosage cup to accurately measure dosage as noted below
- drink plenty of clear fluids to help prevent dehydration, which may accompany diarrhea
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and older	4 teaspoonfuls (1 dosage cup) after the first loose bowel movement; 2 teaspoonfuls (1/2 dosage cup) after each subsequent loose bowel movement; but no more than 8 teaspoonfuls a day
children 9-11 years (60-95 lbs)	2 teaspoonfuls (1/2 dosage cup) after the first loose bowel movement; 1 teaspoonful (1/4 dosage cup) after each subsequent loose bowel movement; but no more than 6 teaspoonfuls a day
children 6-8 years (48-59 lbs)	2 teaspoonfuls (1/2 dosage cup) after the first loose bowel movement; 1 teaspoonful (1/4 dosage cup) after each subsequent loose bowel movement; but no more than 4 teaspoonfuls a day
children under 6 years (up to 47 lbs)	ask a doctor (not intended for use in children under 6 years old)

- avoid excess heat
- store at room temperature, 15° - 30°C (59° - 86°F)



NDC 50383-618-04

Loperamide Hydrochloride Oral Solution

1 mg / 5 mL

Anti-Diarrheal

For the control of the symptoms of diarrhea, including Travelers' Diarrhea.

Each teaspoonful (5 mL) contains:
Loperamide Hydrochloride 1 mg.

Contains Alcohol 0.5%

4 fl oz (118 mL)

Tamper Evident: Do not use if printed band around cap is broken or missing.

Drug Facts

Active Ingredient (in each 5 mL teaspoonful)	Purpose
Loperamide HCl 1 mg	Antidiarrheal
Uses • controls the symptoms of diarrhea, including Travelers' Diarrhea	
Warnings Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl	
Do not use if you have bloody or black stool	
Ask a doctor before use if you have • high fever (greater than 101° F) • mucus present in your stool • a history of liver disease	
Ask a doctor or pharmacist before use if you are taking antibiotics	
Stop use and ask a doctor if diarrhea lasts for more than 2 days	
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions • use and read dosage cup to accurately measure dosage as noted below • drink plenty of clear fluids to help prevent dehydration, which may accompany diarrhea. • find right dosage on chart. If possible, use weight to dose; otherwise, use age.	

adults and children 12 years and older	4 teaspoonfuls (1 dosage cup) after the first loose bowel movement; 2 teaspoonfuls (½ dosage cup) after each subsequent loose bowel movement; but no more than 8 teaspoonfuls a day	ask a doctor (not intended for use in children under 6 years old)
children 9-11 years (60-95 lbs)	2 teaspoonfuls (½ dosage cup) after the first loose bowel movement; 1 teaspoonful (½ dosage cup) after each subsequent loose bowel movement; but no more than 6 teaspoonfuls a day	
children 6-8 years (48-58 lbs)	2 teaspoonfuls (½ dosage cup) after the first loose bowel movement; 1 teaspoonful (½ dosage cup) after each subsequent loose bowel movement; but no more than 4 teaspoonfuls a day	
children under 6 years (up to 47 lbs)		

Other Information • avoid excess heat • See Box for Complete Product Information • store at room temperature, 15°-30°C (59°-86°F)

HI-TECH PHARMACAL CO., INC., Antisville, NY 11701



3 50383-618-04

1 Rev. 6/18/02 12/08

RxCHOICE

Loperamide Hydrochloride Oral Solution
1 mg / 5 mL
Anti-Diarrheal

Drug Facts

Active ingredient

(In each 5 mL, loperamide HCl)

Purpose

Loperamide HCl 1 mg _____ Anti-Diarrheal

Use

• controls the symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Do not use if you have blood or black stool

Ask a doctor before use if you have

• high fever (greater than 101°F)

• recent travel (your stool)

• a history of liver disease

Ask a doctor or pharmacist before use if you are taking antibiotics

Stop use and ask doctor if diarrhea lasts for more than 2 days

If pregnant or breast feeding, ask a health professional before use

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

• use enclosed dosage cup to accurately measure dosage as noted below

• drink plenty of clear fluids to help prevent dehydration, which may accompany diarrhea

• find right dose quickly. If possible, use weight to dose; otherwise, use age.

Drug Facts (continued)

adults and children 12 years and older

4 teaspoonfuls (2 dosage cups) after the first loose bowel movement; 2 teaspoonfuls (1 dosage cup) after each subsequent loose bowel movement, but no more than 8 teaspoonfuls a day

children 6-11 years (60-95 lbs)

2 teaspoonfuls (1 dosage cup) after the first loose bowel movement; 1 teaspoonful (1/2 dosage cup) after each subsequent loose bowel movement, but no more than 4 teaspoonfuls a day

children 6-11 years (30-45 lbs)

2 teaspoonfuls (1 dosage cup) after the first loose bowel movement; 1 teaspoonful (1/2 dosage cup) after each subsequent loose bowel movement, but no more than 4 teaspoonfuls a day

children 6-11 years (15-30 lbs)

ask a doctor (not intended for use in children under 6 years old)

Other information - avoid excess heat

• store at room temperature, 15°-30°C (59°-86°F)

Inactive ingredients: alcohol (0.5%), benzoic acid, cherry flavor, glycerin, mint flavor, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucrose. Citric acid may be used to adjust pH

Questions or comments?

Call 1-800-262-8811

Mon - Thurs 9:00 am - 4:30 pm EST

Fri, Sat 9:00 am - 2:30 pm EST

Get more information associated with use of this product may be reported to this number.

RxCHOICE

NDC 50383-618-04

Loperamide Hydrochloride Oral Solution
1 mg / 5 mL
Anti-Diarrheal

Controls the Symptoms of Diarrhea, including Travelers' Diarrhea

Each 5mL (1 tsp) contains:
Loperamide Hydrochloride 1 mg
contains Alcohol 0.5%

Convenient dosage cup enclosed.

4 fl oz (118 mL)
CHERRY MINT FLAVOR

WATTECH PHARMACAL CO., INC.
Amityville, NY 11701

3 50383-618-04 1

RxCHOICE

Loperamide Hydrochloride Oral Solution
1 mg / 5 mL
Anti-Diarrheal

Controls the Symptoms of Diarrhea, including Travelers' Diarrhea

Each 5mL (1 tsp) contains:
Loperamide Hydrochloride 1 mg
contains Alcohol 0.5%

Convenient dosage cup enclosed.

4 fl oz (118 mL)
CHERRY MINT FLAVOR

WATTECH PHARMACAL CO., INC.
Amityville, NY 11701

3 50383-618-04 1

Inactive ingredients

alcohol (0.5%), benzoic acid, cherry flavor, glycerin, mint flavor, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucrose. Citric acid may be used to adjust pH